

Supplementary Table 2. Univariate and multivariate Cox regression analyses of success in patients with pseudophakia (n = 36)

Variable	Complete success (20%)			Qualified success (20%)		
	HR	95% CI	<i>p</i> -value	HR	95% CI	<i>p</i> -value
Bleb height at 1 mon						
Model 1	1.024	0.545–1.926	0.941	0.765	0.291–2.007	0.586
Model 2	1.071	0.544–2.108	0.842	0.832	0.302–2.291	0.722
Model 3	1.211	0.561–2.616	0.626	0.712	0.174–2.915	0.637
Bleb extent at 1 mon						
Model 1	0.582	0.290–1.168	0.128	0.692	0.227–2.094	0.515
Model 2	0.533	0.241–1.181	0.121	0.713	0.201–2.533	0.601
Model 3	0.554	0.278–1.102	0.092	0.855	0.239–3.054	0.809
Bleb height at 3 mon						
Model 1	0.747	0.366–1.522	0.422	0.757	0.286–2.002	0.574
Model 2	0.713	0.309–1.646	0.428	0.639	0.190–2.152	0.470
Model 3	0.916	0.365–2.302	0.852	0.804	0.221–2.921	0.740
Bleb extent at 3 mon						
Model 1	0.631	0.320–1.244	0.183	0.443	0.153–1.286	0.134
Model 2	0.558	0.268–1.159	0.118	0.372	0.103–1.343	0.131
Model 3	0.446	0.233–0.852	0.015*	0.470	0.135–1.642	0.237
Bleb height at 6 mon						
Model 1	0.657	0.380–1.136	0.133	0.527	0.249–1.118	0.095
Model 2	0.588	0.320–1.081	0.088	0.404	0.164–0.999	0.049*
Model 3	0.906	0.403–2.035	0.811	0.975	0.243–3.914	0.971
Bleb extent at 6 mon						
Model 1	0.627	0.348–1.133	0.122	0.543	0.212–1.390	0.203
Model 2	0.625	0.342–1.142	0.126	0.438	0.154–1.247	0.122
Model 3	0.590	0.308–1.130	0.112	0.388	0.106–1.421	0.153

Model 1, unadjusted univariate analysis; Model 2, adjusted for only age and sex; Model 3, adjusted for age, sex, previous operation history, and diagnosis.

HR = hazard ratio; CI = confidence interval.

*Statistically significant.